Reservoir Storage Outlook


June 18, 2015




DNRC Water Resources Division State Water Projects Bureau



Reservoir Contents Report May 21, 2015



Reservoir Contents Report June 18, 2015



MONTANA DEPARTMENT OF NATURAL RESOURCES AND CONSERVATION

WATER RESOURCES DIVISION - STATE WATER PROJECTS BUREAU

May 31, 2015

All Contents in Acre-Feet

RESERVOIR	TOTAL CAPACITY (includes dead storage)*	CONTENTS								
	Full Pool	AVERAGE	Last Year	Last Month	PRESENT	% CAPACITY	%AVERAGE	READING	COMMENTS	
	Contents	1960 - 2014	5/31/2014	4/30/2015	5/31/2015	5/31/2015	5/31/2015	DATE		
ACKLEY	6,722	4,724	4,552	4,873	6,572	98	139	5/29/2015	elev.= 4317.4	
BAIR	7,300	5,747	7,261	6,370	7,537	103	131	5/28/2015	elev.=5325.84	
COONEY	28,230	24,023	22,584	23,580	28,499	101	119	6/2/2015	elev.=4251.3 (28,409 AF)	
COTTONWOOD	1,900	1,580	1,987	1,900	1,940	102	123	6/2/2015	elev.= 5102.64	
DEADMAN'S BASIN	75,968	54,232	74,686	75,744	75,315	99	139	5/29/2015	elev.=3920.7 (71,565 AF)	
E.F. ROCK CREEK	16,040	10,794	12,005	11,868	10,813	67	100	5/29/2015	elev.=6041.01	
FRENCHMAN	2,777	2,656	2,777	2,777	2,777	100	105	6/1/2015	spilling	
MARTINSDALE	23,348	16,051	20,186	22,924	23,168	99	144	5/29/2015	elev.=4779.1	
MIDDLE CREEK	10,184	9,323	7,935	7,501	10,142	100	109	5/29/2015	elev.=6720.8	
NEVADA CREEK	11,207	11,022	11,522	11,244	9,924	89	90	5/28/2015	elev.=4612.41	
NILAN	10,992	8,531	10,662	10,571	10,980	100	129	6/1/2015	elev.=4442.48	
N.FK. SMITH RIVER	11,406	10,339	11,553	10,399	11,553	101	112	5/28/2015	elev.= 5488.77	
RUBY RIVER	37,612	37,360	37,612	37,844	37,642	100	101	5/5/2015	elev.=5393.0	
TONGUE RIVER	79,071	71,399	83,489	66,537	83,412	105	117	5/29/2015	elev.=3429.5	
W.F. BITTERROOT	32,362	32,617	34,305	32,362	33,334	103	102	5/29/2015	elev.=4726.89	
WILLOW CREEK	18,000	16,618	16,725	16,843	18,809	104	113	6/3/2015	elev.=4736.77	
YELLOWATER	3,842	1,531	3,366	3,187	3,252	85	212	6/1/2015	elev.=3116.95	

^{*} Note: Reservoir contents include dead storage at the following:

Ackley 1001 AF ** O&M slope storage table does not include dead storage (so dead storage has to be added into the storage from the table)

Cooney 90 AF ** Tongue River 711 AF (O&M storage table includes dead storage)

Deadman's 3750 AF ** W. F. Bitterroot 656 AF (O&M storage table includes dead storage)

Nilan 900 AF ** Willow Creek 269 AF (O&M storage table includes dead storage)

^{*} Note: Cooney capacity reflects capacity after 1982 dam rehabilitation; prior capacity was 24,195 A.F.. Average storage shown is for post rehabilitation data.


^{*} Note: Middle Creek capacity reflects capacity after 1993 dam rehabilitation; prior capacity was 8,027 A.F.. Average storage shown is for post rehabilitation data.

^{*} Note: Nevada Creek Reservoir Capacity reflects live storage capacity survey conducted in year 2000. Prior live storage capacity documented as 12,723 AF.

^{*} Note: Tongue River capacity reflects capacity after 1999 dam rehabilitation; prior capacity was 68,040 A.F.. Average storage is post rehabilitation data.

^{*} Note: Frenchman Reservoir capacity tables updated based on aerial survey; prior capacity was 3752 A.F. Average shown is pre aerial survey






- •101% Capacity
- •112% average
- •11,553 Acre-Feet









EMERGENCY SPILLWAY CREST 4780.25

PRINCIPAL SPILLWAY CREST

TRANSDUCER CASE DEPTH

24,350

23,348

8.444

4759.78




- •98% Capacity
- •143% average
- •22,888 Acre-Feet
- •Inflows=0 cfs
- •Outflows=0 cfs

*** PROVISIONAL DATA SUBJECT TO REVISION ***

22,888 AF


*NOTE: RESERVOIR ELEVATIONS BELOW 4759.78 FT ARE NOT VALID DUE TO INSTRUMENTATION

RESERVOIR VOLUME














- •98% Capacity
- •138% average
- •74,674 Acre-Feet (Total Storage)
- •Elev.=3920.4














- •100% Capacity
- •109% Average
- •Outflows~121 cfs
- •10,178 Acre-Feet
- •Elev.=6721.0













- •92% Capacity
- •93% average
- •34,682 Acre-Feet
- •Elev.=5390.0
- •Inflows=211 cfs
- •Outflows=377 cfs


*Data for Ruby Dam is preliminary. Presented data may not accurately represent the actual conditions.













- •101% Capacity
- •119% average
- •28,499 Acre-Feet
- •Elev.=4251.3
- •Inflows= 325 cfs
- •Outflows=325 cfs












- •104% Capacity
- •115% Average
- •82,343 Acre-Feet
- •Elev.=3429.2
- •Inflows=2,310 cfs
- •Outflows=2,180 cfs











1E OF LAST READING	6/17/2015 5:00:00 AM	REFERENCE INFORMATION	FT (MSL)	AC
SERVOIR ELEVATION	6,049.0 FT	DAM CREST	6065.6	19,
SERVOIR VOLUME	13,557 AF	PRINCIPAL SPILLWAY CREST	6055.5	16,
		LOWEST USABLE ELEVATION	5989.7	



- •85% Capacity
- •126% average
- •13,577 Acre-Feet
- •Elev.=6049.0

*** PROVISIONAL DATA SUBJECT TO REVISION ***






,850








- •103% Capacity
- •102 % average
- •33,334 Acre-Feet
- •Elev.=4726.89
- •Inflows~317 cfs
- •Outflows=317 cfs







Summary

- Maximum Winter Carryover was maintained across majority of State Water Projects.
- Reservoirs filled early in spring ahead of normal timing due in part to winter carryover and early snowpack runoff.
- Majority of State Water Projects Reservoirs have average to above average storage for June, Ruby River Reservoir and Nevada Creek Reservoir are below the historical average storage for June.
- Inflows are steadily declining.
- Reservoir releases increasing as decreed water rights come out of priority.









Young Creek

- 5/1-6/30 25cfs
- 7/01-12/31 5cfs

On June 11, FWP measured flows at 14.14 cfs

- •Notification letters were sent to junior water users.
- •Emphasis on encouraging community drought planning.






Tobacco River

- 6/16-6/30 433 cfs
- 7/01-7/15 282 cfs
- •7/16-12/31 100 cfs





Daily discharge, cubic feet per second -- statistics for Jun 17 based on 56 years of record more


Most Recent Instantaneous Value Jun 17					75th percen- tile	
197	201	497	748	784	1030	2270




lowest-

percentile

hydrologic drought




Last updated: 2015-06-17






Map Tools ▼ Basemaps ▼











The second second	Most Recent Instantaneous Value Jun 17	percen-			75th percen- tile	
991	1580	2880	4210	4910	6330	13600






Daily discharge, cubic feet per second -- statistics for Jun 17 based on 30 years of record more

Most Recent Instantaneous Value Jun 17		25th percen- tile			75th percen- tile	
2680	2780	4580	7650	8320	10200	20500








Jun Jun 17 18 2015 2015 Provisional Data Subject to Revision ----Daily discharge, cubic feet per second -- statistics for Jun 18

Most Recent Instantaneous Value Jun 18		25th percen- tile			75th percen- tile	
unavailable	2420	7590	11800	12700	17600	26900

















Discharge, cubic feet per second



△ Median daily statistic (28 years) — Discharge



Montana Outdoors

How to Stay Informed



celebration of Father's Day weekend, June 20-21.



» Learn More



Restrictions & Closures

Submit Public Comments

Contact Us

» Montana WILD Calendar



- Keeping a close eye on streamflows in west and northwest Montana. Plan on engaging with water users to encourage the development of community drought plans.
- FWP will be cooperating with watershed in administering their drought management plans on the Big Hole, Jefferson and Blackfoot Rivers.
- Monitoring temperatures, thermographs on key stream reaches should be completely deployed by July 1. FWP's policy for initiating fishing restrictions and closures is in ARM 12.5.501-509.
- Paddlefish take was delayed due to low streamflows this spring, but conditions did improve and anglers harvested 934 fish.



























SIGNIFICANT FIRE POTENTIAL Valid For: Tuesday, June 23, 2015


Issued On: Wednesday, June 17, 2015 12:33 PM (MT)







NORTHERN ROCKIES GEOGRAPHIC AREA PREDICTIVE SERVICE AREAS








Libby Ranger Station Troy Ranger Station Eureka Ranger Station























West Glacier Hungry Horse Cyclone Benchmark Condon Work Center













Lincoln Phillipsburg





















St. Mary Gleason














West Central Montana




















Northern Plains and Missouri Breaks














Southern Montana (Big Horn/Powder River)

Wolf Mountain Bighorn Mountain Fort Howes Pryor Mountain Badger Peak























— 3% (7)

Current (14)Min

05/27/15

06/10/15

07/08/15

07/22/15

Date

09/02/15

06/24/15

10/28/15

Average

2012

04/15/15

04/29/15

04/01/15

10/14/15

09/30/15

10/28/15

— 3% (10) — Current (18)

> Min Average

2012

04/29/15

05/27/15

07/22/15


Date

06/24/15


04/01/15











National GACC Portal

NRCC Home

About Us Site Disclaimer Contact Us

Wednesday, May 22, 2013

INCIDENT INFORMATION

PREDICTIVE SERVICES

Intelligence

Weather

Fuels/Fire Danger

Outlooks

LOGISTICS / DISPATCH

Dispatch Operations

Aviation

Crews

Equipment/Supplies

Overhead

ADMINISTRATIVE

Northern Rockies Coordinating Group

Policy and Reports

Incident Business Management

Safety Management

Software Applications

Training

RELATED LINKS

National

Area

Welcome to the

NORTHERN ROCKIES COORDINATION CENTER

The Northern Rockies Coordination Center (NRCC) is the interagency focal point for coordinating the mobilization of resources for wildland fire and other all-hazard incidents throughout the Northern Rockies Area and, when necessary, for assignment throughout the United States. Located in Missoula, Montana, the Center also provides Intelligence and Predictive Services related products for use by the wildland fire community for purposes of wildland fire and incident management decision-making.

There are five primary components to the NRCC website.

- Incident Information provides general information on large wildland fires, fire restrictions and closures, and other relevant activity throughout the Geographic Area.
- Predictive Services provides operational products and links to incident situation information, maps, resources, current fire weather conditions, forecasts, fuels, fire behavior as well as daily, weekly and monthly fire weather/fire danger outlooks.
- Logistics/Dispatch provides detailed operation and information links for aviation, crews, equipment and overhead, including Incident Management Teams.
- Administrative provides fire and incident management tools and links including policies and reports, business management, safety, software applications, and training.
- Related Links component provides links to related Internet websites within the Northern Rockies Area and nationally.



BULLETIN BOARD 50



SITUATION

PREPAREDNESS LEVELS Northern Rockies PL: 1 National PL: 1

Situation Reports

Year-to-Date & Historical Wildfire Data

· Restrictions & Closures · · ·

SAFETY ALERTS

NRGA Landscape Mortality Safety Alert NRGA Landscape Mortality Pocket Card

Coal Seam Fires Safety

COOPERATING FEDERAL. STATE AND OTHER AGENCIES IN THE NORTHERN ROCKIES AREA













Montana Drought and Water Supply


Status change from May to June 2015 – Assessed 6/10/2015 (All changes one category unless otherwise noted)

	•	
1)	77	ρr
	\mathbf{L}	$oldsymbol{oldsymbol{oldsymbol{ u}}}$


Lincoln	Gallatin	
Flathead (2)	Wheatland	
Sanders	Sweet Grass	
Lake (2)	Golden Valley	
Mineral	Stillwater	
Missoula	Carbon	
Powell (2)	Musselshell	
Ravalli	Yellowstone	
Granite	Big Horn	
Deer Lodge	Rosebud	
Silver Bow	Phillips	
Cascade	Valley	
Chouteau	Daniels	


No Change

Glacier	Blaine	Prairie
Toole	Meagher	McCone
Pondera	Judith Basin	Dawson
Teton	Fergus	Wibaux
Lewis and	Petroleum	Sheridan
Clark	Park	Richland
Jefferson	Garfield	Roosevelt
Broadwater	Treasure	
Beaverhead	Fallon	
Madison	Powder River	
Liberty	Carter	
Hill	Custer	



Montana Drought Status June 2015 vs. May 2015








Montana Drought & Water Supply Advisory Committee

June 18, 2015 National Weather Service Gina Loss – Service Hydrologist




Departure from Average Temperature May 2015




- West and southwest... near to above average
- Central and east... near to below average




Percent of Normal Precipitation May 2015




May Rankings 53th coolest, 66th driest








Percent of Normal Precipitation Water Year 2015






Percent of Normal Precipitation
Crop Year



Temperature Anomalies June 1 - 15




Highs

- Widespread 3-9 °F above average
- Isolated areas west 9-15 °F above average


• Lows

- Mostly near to 6 °F above average
- South-central to southeast6-10 °F above average





Percent of Average Precipitation June 1 - 16



- Well below average west, north central and portions of south-central and east
- Above average central and east




Flooding in Eastern Montana



 Several Flood Warnings and Flash Flood Warnings were issued for areas from Sidney into south-central and southeast Montana




June 10 - Road is flooded north of Shepherd after a storm dumped up to three inches of rain in the area. Photo: Jeff Branson - Billings Gazette






National Drought Monitor Issued June 18




- D0 'Abnormally Dry' expanded across west southwest and small portion of hi-line
- D1 'Moderate Drought' continues in far southwest and introduced west of the divide
- D0 'Abnormally Dry' reduced over eastern Montana






North American Drought Monitor






Statewide Average Precipitation

June wettest month of year, July drops to nearly half of June






7-Day Precipitation Forecast





8 to 14 Day Outlook 500mb Heights and Anomalies




- June 25 July 1
- Subtle high pressure ridge over Pacific Northwest and Montana




8 to 14 Day Outlook June 25 – July 1

Temperature



• 33% to 70% chance temperatures will average above normal across all except far eastern Montana

Precipitation




• 40% to 50% chance precipitation will average below normal statewide



El Niño / La Niña

El Niño Advisory - Likely El Niño will continue through winter 2016






NOAA - National Weather Service – Building a Weather Ready Nation

July Outlook


Updated June 18

Temperature

Precipitation



- 33% to 40% chance temperatures will average above normal west
- Equal chances for above, below or near normal central and east



- 33% to 50% chance precipitation will be above average over southern half of Montana
- Equal chances for above, below or near normal north




August – October Outlook Updated June 18

Temperature

THREE-MONTH OUTLOOK


• 33% to 60% chance temperatures will average above normal over most of Montana



- 33% to 40% chance precipitation will average above normal across extreme southeast Montana
- Equal chances for above, below, or near average west, central, and north



Drought Outlook through September Issued June 18



 Those areas currently in drought status (D1 – Moderate Drought) now expected to persist or intensify



In Summary...

- While May and early June brought some storms, overall precipitation is near to below normal.
 - Only isolated areas southwest, south-central, and southeast are averaging above normal
 - Montana Drought/Water Supply and National Drought Monitor reflecting these drier conditions
- Some flooding occurred east and southeast
 - Includes some flash flooding from thunderstorms
- Going into drier summer months
 - While some areas of state still in green-up, other areas are drying quickly
- El Niño continued to strengthened, and now likely it will persist through winter
- Drought Outlook currently showing those areas of drought currently in Montana expected to persist.
 - No indication of significant expansion.






Governor's Drought & Water Supply Advisory Committee

USDA NASS Mountain Region Montana Field Office Thomas Chard II Statistician






U.S. Winter Wheat Condition







U.S. Winter Wheat Progress



Montana Winter Wheat June 1 Production Forecast

Total Production

- > 101.2 million bushels
- up 9.36 million bushels or 10 percent from last year


Yield

- > June 1 yield forecast is 44 bushels per acre
- ➤ Up 3 bushels from previous forecast and a year ago.



June 1, 2015 Winter Wheat Yield Bushels and Change From Previous Month





Crop Weather Report Week Ending June 14, 2015

- Topsoil and subsoil moisture conditions were comparable to a year ago but worse than the five year average.
- Crops progress continues to track ahead of last year and the five year progress.

Topsoil Moisture Week Ending June 14, 2015

	This	Last	Last	5-yr avg.
	week	week	year	avg.
Very short	6	6	5	3
Short	29	28	24	12
Adequate	57	57	59	62
Surplus	8	9	12	23

Adequate and surplus condition is 65 percent. Below the 5 year average and last year.

Subsoil Moisture Week Ending June 14, 2015

	This week	Last week	Last year	5-yr avg.
Very short	6	7	5	5
Short	23	22	23	14
Adequate	60	58	66	64
Surplus	11	13	6	17

Adequate and surplus condition is 71 percent. Below the 5 year average and last year.

Winter Wheat Condition Week Ending June 14, 2015

	Very	Poor	Fair	Good	Excellent
	poor				
This week	2	7	34	37	20
Last week	2	7	35	37	19
Last year	2	7	29	51	11
5-yr avg.	2	7	28	46	17

Good and excellent condition is 57 percent. Below the 5 year average and last year.

Spring Wheat Condition Week Ending June 14, 2015

	Very	Poor	Fair	Good	Excellent
	poor				
This week	3	4	33	51	9
Last week	3	3	32	51	11
Last year	1	3	42	48	6
5-yr avg.	1	3	34	53	9

Good and excellent condition is 60 percent. Slightly below the 5 year average, but above last year.

Barley Condition Week Ending June 14, 2015

	Very	Poor	Fair	Good	Excellent
	poor				
This week	1	4	33	43	19
Last week	1	4	33	44	18
Last year	0	4	48	39	9
5-yr avg.	0	4	33	48	15

Good and excellent condition is 62 percent. Slightly below the 5 year average, but above last year.

Dry Peas Condition Week Ending June 14, 2015

	Very	Poor	Fair	Good	Excellent
	poor				
This week	2	3	35	33	14
Last week	2	3	35	31	14
Last year	1	2	36	54	7
5-yr avg.	1	2	22	65	10

Good and excellent condition is 47 percent. Below the 5 year average and last year.

Seeding Completed Week Ending June 14, 2015

	This	Last	Last	5-yr
	week	week	year	avg.
Dry Beans	95	88	85	90
Potatoes	95	89	79	91

Blooming Week Ending June 14, 2015

	This	Last	Last	5-yr
	week	week	year	avg.
Dry Peas	16	7	26	12
Canola	43	19	10	10
Flaxseed	8	0	2	3
Lentils	24	5	16	7

Boot Stage & HeadingWeek Ending June 14, 2015

	This week	Last week	Last year	5-yr avg.
Barley	65	29	7	16
Oats	41	17	16	14
Spring Wheat	42	17	7	10
Winter Wheat	95	90	47	61
Headed Winter Wheat	63	38	21	18
Barley	3	NA	0	1

Livestock Grazing Week Ending June 14, 2015

- 96 percent of Cattle and Calves have been moved to summer ranges, ahead of 90 last year and the five-year average of 88 percent.
- 96 percent of Sheep and Lambs have been moved to summer ranges, ahead of last years 90 percent and the five-year average of 86 percent.

Range & Pasture Feed Condition Week Ending June 14, 2015

	Very	Poor	Fair	Good	Excellent
	poor				
This week	4	16	41	32	7
Last week	4	17	40	32	7
Last year	4	12	31	43	10
5-yr avg.	3	10	27	43	17

Good and excellent condition is 39 percent. Below the 5 year average and last year.

Summary Week ending June 14, 2015

- Soil moistures were below the five year averages
- 6.2 days were suitable for field work during the week, compared to 5.4 days last year and the five year average of 4.3 days
- Planting is almost complete
- Winter wheat and barley has started to head
- Ranchers are wrapping up annual movement of livestock to summer ranges

June Releases

June Hog Report released on June 26

 June Acreage, and June Stocks Reports released on June 29

USDA, NASS, Mountain Region: Montana Field Office

Thomas Chard II, Statistician

1-800-835-2612 or 406-441-1240

Email: nass-mt@nass.usda.gov

Montana Data available at following address:

http://www.nass.usda.gov/Statistics_by_State/Montana/index.asp





Lucas Zukiewicz

Water Supply Specialist (Snow Hydrologist)

USDA-NRCS

Montana Snow Surveys

Lucas.Zukiewicz@mt.usda.gov

406-587-6843

http://www.nrcs.usda.gov/wps/portal/nrcs/main/mt/snow/

Streamflows in the coming months will rely on summer precipitation and groundwater contributions, almost all elevations have melted out, and high elevations are ahead of schedule reducing summer flows.


Photo: Looking towards the headwaters of the Ruby River from the Gravelly Range, MT.

June 17th, 2015. Lucas Zukiewicz




Precipitation























Snowpack




































































Water Supply











Summary

- Monthly mountain precipitation was near to well above normal during the month of May east of the Divide, and near record low west of the Divide.
- June precipitation has been slightly below to below normal for this date at most mountain locations, with a few locations in SW MT receiving above average precipitation.
- Snowpack reached maximum accumulation and began melting earlier than normal this year, and peak basin-wide snow water equivalent was near record low in many basins.
- Rivers in Montana have seen their peak snowmelt driven flows. Elevations with SNOTEL sites in the state have melted out at this point. Zero (of 131) SNOTEL sites have snow remaining on June 18th. Many melt out dates are new record early dates.
- Summer streamflow volumes will be dependent on precipitation, snowpack component of runoff has passed, upper elevations that provide recession flows have melted well ahead of schedule.





Lucas Zukiewicz

Water Supply Specialist (Snow Hydrologist)

USDA-NRCS

Montana Snow Surveys

Lucas.Zukiewicz@mt.usda.gov

406-587-6843

http://www.nrcs.usda.gov/wps/portal/nrcs/main/mt/snow/


Streamflows in the coming months will rely on summer precipitation and groundwater contributions, almost all elevations have melted out, and high elevations are ahead of schedule reducing summer flows.

Photo: Looking towards the headwaters of the Ruby River from the Gravelly Range, MT. June 17th, 2015. Lucas Zukiewicz



DAILY STREAMFLOW CONDITIONS


Mednesday, June 17, 2015 09:30ET



Minimum Discharge for June 18

- 06078500 North Fork Sun River near Augusta
- 12301300 Tobacco River near Eureka
- 12302055 Fisher River near Libby
- 12352500 Bitterroot River near Missoula
- 12354000 St. Regis River near St. Regis
- 12390700 Prospect Creek at Thompson Falls


USGS 06036650 Jefferson River near Three Forks MT (Drainage Area: 9532 square miles, Length of Record: 36 years)



USGS 06036650 Jefferson River near Three Forks MT (Drainage area: 9532 square miles, Length of Record: 36 year) 120000 Cumulative flow between daily 25th and 75th percentiles feet Cumulative streamflow of daily median cubic Lowest observed cumulative flow (2004) 100000 Highest observed cumulative flow (1984) t Observed cumulative flow (2015) in millions 80000 60000 Cumulative Streamflow, 40000 20000 OCT NOV DEC JAN **FEB** JUN JUL SEP MAR **APR** MAY AUG **■USGS** WaterWatch 2015

Last updated: 2015-06-17

USGS 06037500 Madison River near West Yellowstone MT (Drainage Area: 420 square miles, Length of Record: 101 years)




Explanation - Percentile classes						
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow	
Much below normal	Below normal	Normal	Above normal	Much above normal		

(Drainage area: 420 square miles, Length of Record: 87 year) 25000 Cumulative flow between daily 25th and 75th percentiles cubic feet Cumulative streamflow of daily median Lowest observed cumulative flow (1934) 20000 Highest observed cumulative flow (1997) 4 Observed cumulative flow (2015) Cumulative Streamflow, in millions 15000 10000 5000 OCT NOV DEC JAN **FEB** MAR **APR** MAY JUN JUL AUG SEP **■USGS** WaterWatch 2015 Last updated: 2015-06-17


USGS 06037500 Madison River near West Yellowstone MT

USGS 06043500 Gallatin River near Gallatin Gateway MT (Drainage Area: 825 square miles, Length of Record: 125 years)




Explanation - Percentile classes						
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow	
Much below normal	Below normal	Normal	Above normal	Much above normal		

USGS 06043500 Gallatin River near Gallatin Gateway MT (Drainage area: 825 square miles, Length of Record: 84 year)




USGS 06054500 Missouri River at Toston MT (Drainage Area: 14669 square miles, Length of Record: 125 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 06054500 Missouri River at Toston MT (Drainage area: 14669 square miles, Length of Record: 81 year)




USGS 06078500 North Fork Sun River near Augusta MT (Drainage Area: 258 square miles, Length of Record: 103 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 06099500 Marias River near Shelby MT (Drainage Area: 3242 square miles, Length of Record: 112 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 06115200 Missouri River near Landusky MT (Drainage Area: 40987 square miles, Length of Record: 80 years)




USGS 06120500 Musselshell River at Harlowton MT (Drainage Area: 1125 square miles, Length of Record: 107 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 06191500 Yellowstone River at Corwin Springs MT (Drainage Area: 2619 square miles, Length of Record: 125 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 06191500 Yellowstone River at Corwin Springs MT (Drainage area: 2619 square miles, Length of Record: 108 year)




USGS 06200000 Boulder River at Big Timber MT (Drainage Area: 523 square miles, Length of Record: 67 years)






USGS 06200000 Boulder River at Big Timber MT (Drainage area: 523 square miles, Length of Record: 65 year)




USGS 06214500 Yellowstone River at Billings MT (Drainage Area: 11805 square miles, Length of Record: 86 years)






USGS 06214500 Yellowstone River at Billings MT (Drainage area: 11805 square miles, Length of Record: 86 year)




USGS 06287000 Bighorn River near St. Xavier, MT (Drainage Area: 19667 square miles, Length of Record: 80 years)




E	xplana	tion - Pe	ercentile	classes	
lowest- 10th percentile	10-24	25-75	76-90	90th percentile	Flow
Much below normal	Below normal	Normal	Above	-highest Much above normal	FIOW

USGS 06287000 Bighorn River near St. Xavier, MT (Drainage area: 19667 square miles, Length of Record: 80 year)




USGS 06306300 Tongue River at State Line nr Decker MT (Drainage Area: 1453 square miles, Length of Record: 54 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 06306300 Tongue River at State Line nr Decker MT (Drainage area: 1453 square miles, Length of Record: 54 year)




USGS 06324500 Powder River at Moorhead MT (Drainage Area: 8086 square miles, Length of Record: 85 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 06329500 Yellowstone River near Sidney MT (Drainage Area: 69083 square miles, Length of Record: 104 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 12301300 Tobacco River near Eureka MT (Drainage Area: 440 square miles, Length of Record: 56 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 12302055 Fisher River near Libby MT (Drainage Area: 838 square miles, Length of Record: 47 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 12302055 Fisher River near Libby MT (Drainage area: 838 square miles, Length of Record: 47 year)




USGS 12340000 Blackfoot River near Bonner MT (Drainage Area: 2290 square miles, Length of Record: 116 years)






USGS 12340000 Blackfoot River near Bonner MT (Drainage area: 2290 square miles, Length of Record: 79 year)




USGS 12340500 Clark Fork above Missoula MT (Drainage Area: 5999 square miles, Length of Record: 85 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below	Below normal	Normal	Above	Much above			

USGS 12340500 Clark Fork above Missoula MT (Drainage area: 5999 square miles, Length of Record: 85 year)




USGS 12344000 Bitterroot River near Darby MT (Drainage Area: 1049 square miles, Length of Record: 77 years)




Explanation - Percentile classes						
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow	
Much below normal	Below normal	Normal	Above normal	Much above normal		

USGS 12344000 Bitterroot River near Darby MT (Drainage area: 1049 square miles, Length of Record: 77 year)




USGS 12352500 Bitterroot River near Missoula MT (Drainage Area: 2814 square miles, Length of Record: 116 years)




Explanation - Percentile classes						
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow	
Much below normal	Below normal	Normal	Above normal	Much above normal		

USGS 12354000 St. Regis River near St. Regis, MT (Drainage Area: 303 square miles, Length of Record: 104 years)




Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

USGS 12354500 Clark Fork at St. Regis MT (Drainage Area: 10709 square miles, Length of Record: 85 years)




Explanation - Percentile classes						
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow	
Much below normal	Below normal	Normal	Above normal	Much above		

USGS 12354500 Clark Fork at St. Regis MT (Drainage area: 10709 square miles, Length of Record: 85 year)




USGS 12358500 M F Flathead River near West Glacier MT (Drainage Area: 1128 square miles, Length of Record: 75 years)



Explanation - Percentile classes						
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow	
Much below normal	Below normal	Normal	Albove normal	Much above normal		


USGS 12358500 M F Flathead River near West Glacier MT (Drainage area: 1128 square miles, Length of Record: 75 year) Cumulative flow between daily 25th and 75th percentiles Cumulative streamflow of daily median Lowest observed cumulative flow (1941) Highest observed cumulative flow (2011) Observed cumulative flow (2015)



140000


■USGS WaterWatch

USGS 12389000 Clark Fork near Plains MT (Drainage Area: 19958 square miles, Length of Record: 104 years)




Explanation - Percentile classes						
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow	
Much below normal	Below normal	Normal	Above normal	Much above normal		

USGS 12389000 Clark Fork near Plains MT (Drainage area: 19958 square miles, Length of Record: 104 year)



USGS 12390700 Prospect Creek at Thompson Falls MT (Drainage Area: 182 square miles, Length of Record: 58 years)



Explanation - Percentile classes							
lowest- 10th percentile	10-24	25-75	76-90	90th percentile -highest	Flow		
Much below normal	Below normal	Normal	Above normal	Much above normal			

